

## Are you ready for Calculus? Solutions

$$1. a. \quad \frac{x(x^2 - 9)}{(x-4)(x-3)} = \frac{x(x-3)(x+3)}{(x-4)(x-3)} = \frac{x(x+3)}{x-4}$$

$$1. b. \quad \frac{x^2 - 2x - 8}{x^3 + x^2 - 2x} = \frac{(x-4)(x+2)}{x(x^2 + x - 2)} = \frac{(x-4)(x+2)}{x(x-1)(x+2)} = \frac{x-4}{x(x-1)}$$

$$1. c. \quad \left[ \frac{\frac{1}{x} - \frac{1}{5}}{\frac{1}{x^2} - \frac{1}{25}} \right] \frac{25x^2}{25x^2} = \frac{(25x - 5x^2)}{(25 - x^2)} = \frac{5x(5-x)}{(5-x)(5+x)} = \frac{5x}{5+x}$$

$$1. d. \quad \left[ \frac{9 - \frac{1}{x^2}}{3 + \frac{1}{x}} \right] \frac{x^2}{x^2} = \frac{9x^2 - 1}{3x^2 + x} = \frac{(3x-1)(3x+1)}{x(3x+1)} = \frac{3x-1}{x}$$

$$2. a. \quad \left( \frac{2}{\sqrt{3} + \sqrt{2}} \right) \left( \frac{\sqrt{3} - \sqrt{2}}{\sqrt{3} - \sqrt{2}} \right) = \frac{2(\sqrt{3} - \sqrt{2})}{3-2} = 2(\sqrt{3} - \sqrt{2})$$

$$2. b. \quad \left( \frac{4}{1 - \sqrt{5}} \right) \left( \frac{1 + \sqrt{5}}{1 + \sqrt{5}} \right) = \frac{4(1 + \sqrt{5})}{1 + \sqrt{5} - \sqrt{5} - 5} = \frac{4(1 + \sqrt{5})}{-4} = -(1 + \sqrt{5}) = -1 - \sqrt{5}$$

$$2. c. \quad \left( \frac{1}{(1 + \sqrt{3}) - \sqrt{5}} \right) \left( \frac{(1 + \sqrt{3}) + \sqrt{5}}{(1 + \sqrt{3}) + \sqrt{5}} \right) = \frac{(1 + \sqrt{3}) + \sqrt{5}}{(1 + \sqrt{3})^2 - (\sqrt{5})^2} = \frac{(1 + \sqrt{3}) + \sqrt{5}}{1 + 2\sqrt{3} + 3 - 5}$$

$$= \left( \frac{1 + \sqrt{3} + \sqrt{5}}{-1 + 2\sqrt{3}} \right) \left( \frac{-1 - 2\sqrt{3}}{-1 - 2\sqrt{3}} \right) = \frac{-1 - 2\sqrt{3} - 1\sqrt{3} - 6 - 1\sqrt{5} - 2\sqrt{15}}{1 - 12}$$

$$= \frac{-1 - 2\sqrt{3} - 1\sqrt{3} - 6 - 1\sqrt{5} - 2\sqrt{15}}{1 - 12} = \frac{-7 - 3\sqrt{3} - \sqrt{5} - 2\sqrt{15}}{-11}$$

$$= \frac{7 + 3\sqrt{3} + \sqrt{5} + 2\sqrt{15}}{11}$$

$$3. a. \quad \frac{(2a^2)^3}{b} = \frac{8a^6}{b} \quad [= 8a^6b^{-1}]$$

$$3. b. \quad \sqrt{9ab^3} = (9ab^3)^{\frac{1}{2}} = 3a^{\frac{1}{2}}b^{\frac{3}{2}}$$

$$3. c. \quad \frac{\frac{2a}{b}}{\frac{3}{a}} = \frac{2a}{b} \cdot \frac{a}{3} = \frac{2a^2}{3b} \quad \left[ = \frac{2}{3}a^2b^{-1} \right]$$

$$3. d. \quad \frac{a(b-1)}{b(b-1)} = \frac{a}{b} \quad [= ab^{-1}]$$

$$3. e. \quad \frac{b}{a\sqrt{a}} = \frac{b}{a^{\frac{3}{2}}} \quad \left[ = a^{-\frac{3}{2}}b \right]$$

$$3. f. \quad \frac{a^{\frac{4}{3}}b^{\frac{3}{2}}}{ba^{\frac{1}{2}}} = \frac{a^{\frac{8}{6}}b^{\frac{3}{2}}}{ba^{\frac{3}{6}}} = a^{\frac{5}{6}}b^{\frac{1}{2}}$$

$$4. a. \quad 5^{x+1} = 5^2$$

$$x+1=2$$

$$x=1$$

$$4. b. \quad 3^{-1} = 3^{2x+2}$$

$$2x+2 = -1$$

$$2x = -3$$

$$x = -\frac{3}{2}$$

$$4. c. \quad x = 2^3 = 8$$

$$4. d. \quad \log_3 x^2 = \log_3 4^2 - \log_3 5^4$$

$$\log_3 x^2 = \log_3 \frac{4^2}{5^4}$$

$$x^2 = \frac{4^2}{5^4}$$

$$x = \pm \frac{4}{5^2} = \pm \frac{4}{25}$$

$$5. a. \quad \log_2 \frac{5(x^2-1)}{x-1} = \log_2 \frac{5(x-1)(x+1)}{x-1} = \log_2 5(x+1)$$

$$5. b. \quad \log_4 9^2 - \log_2 3 = \frac{\log_2 9^2}{\log_2 4} - \log_2 3 = \frac{\log_2 9^2}{2} - \log_2 3 = \log_2 (9^2)^{\frac{1}{2}} - \log_2 3$$

$$= \log_2 9 - \log_2 3 = \log_2 \frac{9}{3} = \log_2 3$$

$$5. c. \quad 3^{\log_3 5^2} = 25 \quad 6. a. \quad \frac{1}{2} \quad 6. b. \quad \log_{10} (10^{-x}) = -x$$

$$6. c. \quad \log_{10} (\sqrt{x})^2 + \log_{10} \left(x^{\frac{1}{3}}\right)^3 = \log_{10} x + \log_{10} x = 2\log_{10} x \quad [= \log_{10} x^2]$$

$$7. a. \quad \frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

$$\frac{x}{a} = 1 - \frac{y}{b} - \frac{z}{c}$$

$$x = a \left[ 1 - \frac{y}{b} - \frac{z}{c} \right]$$

$$a = \frac{x}{\left[ 1 - \frac{y}{b} - \frac{z}{c} \right]} \left[ \frac{bc}{bc} \right]$$

$$a = \frac{bcx}{bc - cy - cz}$$

$$7. b. \quad \frac{V}{2} = ab + bc + ca$$

$$\frac{V}{2} - bc = ab + ac$$

$$\frac{V}{2} - bc = a(b+c)$$

$$a = \frac{\left[ \frac{V}{2} - bc \right]}{\left[ \frac{2}{b+c} \right]} \left[ \frac{2}{2} \right]$$

$$a = \frac{V - 2bc}{2(b+c)}$$

$$7.c. \quad A = 2\pi r^2 + 2\pi r h$$

$$0 = 2\pi r^2 + 2\pi r h - A$$

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$a = 2\pi \quad b = 2\pi h, \quad c = -A$$

$$r = \frac{-2\pi h \pm \sqrt{(2\pi h)^2 - 4(2\pi)(-A)}}{2(2\pi)}$$

$$= \frac{-2\pi h \pm \sqrt{4\pi^2 h^2 + 8A\pi}}{4\pi}$$

$$= \frac{-\pi h \pm \sqrt{\pi^2 h^2 + 2A\pi}}{2\pi}$$

only asks for positive r so

$$r = \frac{-\pi h + \sqrt{\pi^2 h^2 + 2A\pi}}{2\pi}$$

$$7.d. \quad A = P + nrP$$

$$A = P(1 + nr)$$

$$P = \frac{A}{1 + nr}$$

$$7.e. \quad 2x - y = xd + 2yd$$

$$2x - y = d(x + 2y)$$

$$d = \frac{2x - y}{x + 2y}$$

$$8.a. \quad y = (x^2 + 4x + 4) + 3 - 4$$

$$y = (x^2 + 4x + 4) + 3 - 4$$

$$y = (x + 2)^2 - 1$$

$$y + 1 = (x + 2)^2$$

$$y - (-1) = (x - (-2))^2$$

$$7.f. \quad \frac{2x}{4\pi} + \frac{1-x}{2} = 0$$

$$2\pi \left[ \frac{x}{2\pi} + \frac{1-x}{2} = 0 \right]$$

$$x + \pi(1-x) = 0$$

$$x + \pi - x\pi = 0$$

$$x - x\pi = -\pi$$

$$x(1 - \pi) = -\pi$$

$$x = \frac{-\pi}{1 - \pi} = \frac{\pi}{\pi - 1}$$

$$8.b. \quad -2y = 3x^2 + 3x$$

$$-2y = 3(x^2 + x + \quad)$$

$$-2y = 3\left(x^2 + x + \frac{1}{4}\right) - \frac{3}{4}$$

$$-2y + \frac{3}{4} = 3\left(x + \frac{1}{2}\right)^2$$

$$-2\left(y - \frac{3}{8}\right) = 3\left(x + \frac{1}{2}\right)^2$$

$$y - \frac{3}{8} = -\frac{3}{2}\left(x - \left(-\frac{1}{2}\right)\right)^2$$

$$8. c. \quad x + 9 = 9 \left( y^2 - \frac{2}{3}y \right)$$

$$x + 9 = 9 \left( y^2 - \frac{2}{3}y + \frac{1}{9} \right) - 1$$

$$x + 10 = 9 \left( y^2 - \frac{2}{3}y + \frac{1}{9} \right)$$

$$x - (-10) = 9 \left( y - \frac{1}{3} \right)^2$$

$$9. a. \quad x^6 - 16x^4 = x^4 (x^2 - 16) = x^4 (x + 4)(x - 4)$$

$$9. b. \quad 4x^3 - 8x^2 - 25x + 50 = 4x^2(x - 2) - 25(x - 2) = (x - 2)(4x^2 - 25) = (x - 2)(2x + 5)(2x - 5)$$

$$9. c. \quad 8x^3 + 27 = (2x + 3)(4x^2 - 6x + 9)$$

$$a^3 + b^3 = (a + b)(a^2 + ab + b^3)$$

$$9. d. \quad x^4 - 1 = (x^2 + 1)(x^2 - 1)$$

$$= (x^2 + 1)(x + 1)(x - 1)$$

$$10. a. \quad x^6 - 16x^4 = 0 \quad x^4 = 0 \quad \text{or} \quad x + 4 = 0 \quad \text{or} \quad x - 4 = 0$$

$$x^4 (x^2 - 16) = 0 \quad x = 0 \quad x = -4 \quad x = 4$$

$$x^4 (x + 4)(x - 4) = 0 \quad 0, \pm 4$$

$$10. b. \quad (x - 2)(2x + 5)(2x - 5) = 0$$

$$x - 2 = 0 \quad \text{or} \quad 2x + 5 = 0 \quad \text{or} \quad 2x - 5 = 0 \quad 2, \pm \frac{5}{2}$$

$$x = 2 \quad x = -\frac{5}{2} \quad x = \frac{5}{2}$$

$$10. c. \quad (2x + 3)(4x^2 - 6x + 9) = 0$$

$$2x + 3 = 0 \quad \text{or} \quad 4x^2 - 6x + 9 = 0$$

$$x = -\frac{3}{2} \quad \sqrt{b^2 - 4ac} = \sqrt{(-6)^2 - 4(4)9} = \sqrt{36 - 72}$$

the square root of a negative is not real

$$11. a. \quad 3 \sin^2 x = \cos^2 x$$

$$\frac{\sin^2 x}{\cos^2 x} = \frac{1}{3}$$

$$\tan^2 x = \frac{1}{3} \quad \Rightarrow \quad \tan x = \pm \frac{1}{\sqrt{3}} = \pm \frac{\sqrt{3}}{3} \quad \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$$

11. b.  $(1 - \sin^2 x) - \sin^2 x = \sin x$

$1 - 2\sin^2 x - \sin x = 0$

$0 = 2\sin^2 x + \sin x - 1$

$0 = (2\sin x - 1)(\sin x + 1)$

$2\sin x - 1 = 0$  or  $\sin x + 1 = 0$  ( $-\pi < x < \pi$ )

$\sin x = \frac{1}{2}$                        $\sin x = -1$

$x = \frac{\pi}{6}, \frac{5\pi}{6}$                        $x \left( = \frac{3\pi}{2} \right) = -\frac{\pi}{2}$

11. c.  $\tan x + \sec x = 2 \cos x$

$\cos x \left[ \frac{\sin x}{\cos x} + \frac{1}{\cos x} = 2 \cos x \right]$

$\sin x + 1 = 2 \cos^2 x$

$\sin x + 1 = 2(1 - \sin^2 x)$

$\sin x + 1 = 2 - 2\sin^2 x$

$2\sin^2 x + \sin x - 1 = 0$







$(2\sin x - 1)(\sin x + 1) = 0$

$2\sin x - 1 = 0$  or  $\sin x + 1 = 0$


$\sin x = \frac{1}{2}$                        $\sin x = -1$

$x = \frac{\pi}{6}, \frac{5\pi}{6}$                        $x = \frac{3\pi}{2}$  (tan x and sec x are undefined here)


$x = \frac{\pi}{6} + 2k\pi$  or  $\frac{5\pi}{6} + 2k\pi$  where k is any integer




13. a.  $\sin\left(x - \frac{\pi}{4}\right)$   $b=1$   $c=\frac{\pi}{4}$  phase shift =  $\frac{c}{b} = \frac{\frac{\pi}{4}}{1} = \frac{\pi}{4}$  to right




13. b.  $\sin\frac{x}{2}$  horizontal stretch 2 times




13. c.  $2\sin x$  magnitude increased 2 times



13. d.  $\cos x$



13. e.  $\frac{1}{\sin x} = \csc x$  Draw  $\sin x$  curve to help draw  $\csc x$  curve,  $-\pi$ ,  $0$ , and  $\pi$  are asymptotes



14. a.  $4x^2 + 12x + 3 = 0$  not factorable - use quadratic formula

$a = 4$   $b = 12$   $c = 3$

$$x = \frac{-12 \pm \sqrt{12^2 - 4(4)3}}{2(4)} = \frac{-12 \pm \sqrt{144 - 48}}{8} = \frac{-12 \pm \sqrt{96}}{8} = \frac{-12 \pm 4\sqrt{6}}{8} = \frac{-3 \pm \sqrt{6}}{2}$$

$$14. b. \quad (2x + 1)(x + 2) = 5$$

$$2x^2 + 4x + x + 2 = 5$$

$$2x^2 + 5x + 2 - 5 = 0$$

$$2x^2 + 5x - 3 = 0$$

$$(2x - 1)(x + 3) = 0$$

$$2x - 1 = 0 \quad \text{or} \quad x + 3 = 0 \quad -3, \frac{1}{2}, (-2 \text{ is not in the domain})$$

$$x = \frac{1}{2} \quad x = -3$$

$$14. c. \quad x(x+1) \left[ \frac{x+1}{x} - \frac{x}{x+1} = 0 \right]$$

$$(x+1)^2 - x^2 = 0$$

$$x^2 + 2x + 1 - x^2 = 0$$

$$2x + 1 = 0$$

$$x = -\frac{1}{2} \quad (x = 0 \text{ and } -1 \text{ are not in the domain})$$

$$15. a. \quad \begin{array}{r} \underline{-2 \mid} \quad 1 \quad -4 \quad 1 \quad 0 \quad -7 \quad 1 \\ \quad \quad -2 \quad 12 \quad -26 \quad 52 \quad -90 \\ \hline 1 \quad -6 \quad 13 \quad -26 \quad 45 \quad -89 \end{array}$$

Remainder -89

$$15. b. \quad \begin{array}{r} x^2 - x + 1 \\ x^3 + 1 \overline{) x^5 - x^4 + x^3 + 2x^2 - x + 4} \\ \underline{x^5} \phantom{+ x^3} + x^2 \\ -x^4 + x^3 + x^2 - x + 4 \\ \underline{-x^4} \phantom{+ x^3} -x \\ \phantom{-x^4} x^3 + x^2 + 4 \\ \underline{x^3} \phantom{+ x^2} + 1 \\ \phantom{-x^4} \phantom{x^3} x^2 + 3 \end{array}$$

Remainder  $x^2 + 3$

$$16. a. \quad \begin{array}{r} \underline{2 \mid} \quad 12 \quad -23 \quad -3 \quad 2 \\ \quad \quad 24 \quad 2 \quad -2 \\ \hline 12 \quad 1 \quad -1 \quad 0 \end{array}$$

$$(12x^2 + x - 1) = (4x - 1)(3x + 1)$$

$$12x^3 - 23x^2 - 3x + 2 = (x - 2)(4x - 1)(3x + 1) \quad \left\{ 2, \frac{1}{4}, -\frac{1}{3} \right\}$$

16. b. Rational zero (root) theorem  $p = \pm 1$   $q = \pm 12, \pm 6, \pm 4, \pm 3, \pm 2, \pm 1$

$$\frac{p}{q} = \pm \frac{1}{12}, \pm \frac{1}{6}, \pm \frac{1}{4}, \pm \frac{1}{3}, \pm \frac{1}{2}, \pm 1$$

$$\begin{array}{r} \cdot \frac{1}{2} \mid \quad 12 \quad 8 \quad -1 \quad -1 \\ \quad \quad \quad \underline{\quad 6 \quad 7 \quad} \\ 12 \quad 14 \quad 6 \end{array} \quad \text{Not a solution}$$

$$\begin{array}{r} \cdot \frac{-1}{2} \mid \quad 12 \quad 8 \quad -1 \quad -1 \\ \quad \quad \quad \underline{\quad -6 \quad -1 \quad 1 \quad} \\ 12 \quad 2 \quad -2 \quad 0 \end{array}$$

$$(12x^2 + 2x - 2) = 2(6x^2 + 1x - 1) = 2(3x - 1)(2x + 1)$$

$$\left\{ -\frac{1}{2}, \frac{1}{3}, -\frac{1}{2} \right\}$$


17. a.  $x^2 + 2x - 3 \leq 0$

$$(x - 1)(x + 3) \leq 0$$

$$x - 1 = 0 \quad \text{or} \quad x + 3 = 0$$

$$x = 1 \quad \quad \quad x = -3$$

zeros -3, 1



$$[-3, 1] \text{ or } -3 \leq x \leq 1$$

17. b. If  $3x - 2 > 0$   $x > \frac{2}{3}$

$$2x - 1 \leq 3x - 2$$


$$1 \leq x$$

If  $3x - 2 < 0$

$$x < \frac{2}{3}$$

~~$$2x - 1 \geq 3x - 2$$~~

~~$$1 \geq x$$~~




$$x \geq 1 \text{ or } x < \frac{2}{3}$$

17. c.  $(x^2 + x + \quad) + 1 > 0$

$$\left(x^2 + x + \frac{1}{4}\right) + 1 - \frac{1}{4} > 0$$

$$\left(x^2 + x + \frac{1}{4}\right) + \frac{3}{4} > 0$$



never less than zero, so all x

18. a.  $-1 \leq -x + 4 \leq 1$

$$\underline{-4 \quad -4 \quad -4}$$

$$-5 \leq -x \leq -3$$

$$5 \geq x \geq 3$$

18. b.  $5x - 2 = 8$

$$\underline{\quad +2 \quad +2}$$

$$5x = 10$$

$$x = 2$$

or  $-(5x - 2) = 8$

$$-5x + 2 = 8$$

$$\underline{\quad -2 \quad -2}$$

$$-5x = 6$$

$$x = -\frac{6}{5}$$



$$18. c. \quad 2x+1 = x+3 \quad \text{or} \quad -(2x+1) = x+3$$

$$x = 2 \quad -2x - 1 = x+3$$

$$-4 = 3x$$

$$x = -\frac{4}{3}$$

$$19. a. \quad m = \frac{-4-3}{2--1} = \frac{-7}{3} \quad y+4 = \frac{-7}{3}(x-2)$$

$$3(y+4) = -7(x-2)$$

$$3y+12 = -7x+14 \quad \text{or} \quad 3y = -7x+2$$

$$7x + 3y = 2 \quad y = -\frac{7}{3}x + \frac{2}{3}$$

$$19. b. \quad 2x - 3y + 5 = 0 \quad \perp m = \frac{-3}{2} \quad y-2 = \frac{-3}{2}(x+1)$$

$$-3y = -2x - 5 \quad 2(y-2) = -3(x+1)$$

$$y = \frac{2}{3}x + \frac{5}{3} \Rightarrow m = \frac{2}{3} \quad 2y - 4 = -3x - 3$$

$$\text{or } m = -\frac{A}{B} = -\frac{2}{-3} = \frac{2}{3} \quad 3x + 2y = 1$$

$$19. c. \quad \left( \frac{-1+3}{2}, \frac{4+2}{2} \right) = (1, 3) \quad y=3$$

$$20. a. \quad y = 3x - 7 \quad \text{Substitute} \quad x + 5(3x - 7) + 3 = 0$$


$$x + 15x - 35 + 3 = 0$$

$$16x - 32 = 0$$

$$x = 2 \quad y = 3(2) - 7 = -1 \quad (2, -1)$$

$$20. b. \quad y \geq 3x - 7 \quad \text{and} \quad 5y \geq -x - 3$$

$$y \geq -\frac{x}{5} - \frac{3}{5}$$



$$21. a. \quad (x-h)^2 + (y-k)^2 = R^2 \quad \text{where } (h, k) \text{ is the center of the circle and } R \text{ is the radius}$$


$$(x-1)^2 + (y-2)^2 = R^2$$

$$(-2-1)^2 + (-1-2)^2 = R^2$$

$$9 + 9 = R^2$$

$$18 = R^2 \Rightarrow (x-1)^2 + (y-2)^2 = 18$$

21. b. Uses a Geometry rule that says the segment perpendicular to the midpoint of a chord must be a diameter of the circle (i. e. go through the center of the circle)



The midpoint of  $(0, 0)$  &  $(0, 2)$  is  $(0, 1)$   
 and the midpoint of  $(0, 0)$  &  $(1, 0)$  is  $(\frac{1}{2}, 0)$   
 so the center of the circle is  $(\frac{1}{2}, 1)$

continued next page

21. b. cont.  $(x-h)^2 + (y-k)^2 = R^2$  where  $(h, k)$  is the center of the circle and  $R$  is the radius

$$\left(x - \frac{1}{2}\right)^2 + (y-1)^2 = R^2 \quad \text{using } (0,0) \text{ as the point on the circle}$$

$$\frac{1}{4} + 1 = R^2$$

$$\frac{5}{4} = R^2 \Rightarrow \left(x - \frac{1}{2}\right)^2 + (y-1)^2 = \frac{5}{4}$$

22. a.  $(x-h)^2 + (y-k)^2 = R^2$  where  $(h, k)$  is the center of the circle and  $R$  is the radius

$$(x^2 + 6x) + (y^2 - 4y) + 3 = 0 \quad \text{Complete the squares}$$

$$(x^2 + 6x + 9) + (y^2 - 4y + 4) + 3 - 9 - 4 = 0$$

$$(x + 3)^2 + (y - 2)^2 - 10 = 0$$

$$(x + 3)^2 + (y - 2)^2 = 10 \Rightarrow \text{center } (-3, 2) \text{ \& Radius } \sqrt{10}$$

b. The slope of the radius from  $(-3, 2)$  to  $(-2, 5)$  is

$$m = \frac{5-2}{-2-(-3)} = \frac{3}{1} = 3 \quad \text{to be tangent to the circle a line must be perpendicular to the radius}$$


so the slope of the tangent line is  $-\frac{1}{3}$

$$(y-5) = -\frac{1}{3}(x+2)$$

$$3y - 15 = -x - 2$$

$$x + 3y = 13$$

23.



The center is on  $y = 3$  or  $k = 3$

two points on the circle is  $(1, 0)$  &  $(0, 3)$

$$(1-h)^2 + (0-3)^2 = R^2$$

$$(0-h)^2 + (3-3)^2 = R^2$$

$$(1-h)^2 + (0-3)^2 = (0-h)^2 + (3-3)^2$$

$$1 - 2h + h^2 + 9 = h^2$$

$$10 - 2h = 0$$

$$10 = 2h$$

$$h = 5$$

b. Center is  $(5, 3)$

$$(0-5)^2 + (3-3)^2 = R^2$$

$$25 = R^2$$

$$(x-5)^2 + (y-3)^2 = 25$$

a. Call the other  $x$ -intercept  $(a, 0)$

$$(a-5)^2 + (0-3)^2 = 25$$

$$a^2 - 10a + 25 + 9 = 25$$

$$a^2 - 10a + 9 = 0$$

$$(a-9)(a-1) = 0 \Rightarrow a = 1 \text{ or } 9 \quad (9, 0)$$

24. Distance from A =  $\sqrt{(x+1)^2 + (y-1)^2}$  and Distance from B =  $\sqrt{(x-2)^2 + (y+1)^2}$

Distance from A = 3(Distance from B)

$$\sqrt{(x+1)^2 + (y-1)^2} = 3\sqrt{(x-2)^2 + (y+1)^2}$$

$$(x+1)^2 + (y-1)^2 = 9((x-2)^2 + (y+1)^2)$$

$$[x^2 + 2x + 1 + y^2 - 2y + 1] = 9[x^2 - 4x + 4 + y^2 + 2y + 1]$$


$$x^2 + 2x + 1 + y^2 - 2y + 1 = 9x^2 - 36x + 36 + 9y^2 + 18y + 9$$

$$0 = 8x^2 + 8y^2 - 38x + 20y + 43$$

25. a.  $x^2 + x - 2 > 0$

$$(x+2)(x-1) > 0$$

Domain  $(-\infty, -2)$  and  $(1, \infty)$



25. b. i. Domain all real numbers, Range 7

25. b. ii. Domain all real numbers except  $x = \frac{1}{2}$  (vertical asymptote),

Range all real numbers except  $\frac{5}{2}$  (horizontal asymptote)

26.  $x > 0$   $f(x) = \frac{x}{x} = 1$        $x < 0$   $f(x) = \frac{x}{-x} = -1$        $x = 0$   $f(x)$  is undefined

Domain all real numbers except 0, Range  $\{-1, 1\}$

27. a.  $\frac{[2(x+h)+3] - [2x+3]}{h} = \frac{2x+2h+3 - 2x-3}{h} = \frac{2h}{h} = 2$

27. b.  $\frac{\left[\frac{1}{x+h+1}\right] - \left[\frac{1}{x+1}\right]}{h} = \frac{(x+h+1)(x+1)}{(x+h+1)(x+1)} \left[ \frac{\left[\frac{1}{x+h+1}\right] - \left[\frac{1}{x+1}\right]}{h} \right] = \frac{(x+1) - (x+h+1)}{h(x+h+1)(x+1)}$


$$= \frac{x+1-x-h-1}{h(x+h+1)(x+1)} = \frac{-h}{h(x+h+1)(x+1)} = \frac{-1}{(x+h+1)(x+1)}$$

27. c.  $\frac{(x+h)^2 - x^2}{h} = \frac{x^2 + 2xh + h^2 - x^2}{h} = \frac{2xh + h^2}{h} = 2x + h$


28. a. Shifts one unit to left




28. b. Reflects across y-axis




28. c. all positive y values



28. d. negative x-axis is reflection of positive x-axis




29. a. Draw  $y = 3x + 2$  – do not go below the x-axis, reflect across  $x = -\frac{2}{3}$



29. b.  $(x^2 + x + \quad)$

$$\left(x^2 + x + \frac{1}{4}\right) + \frac{-1}{4}$$

$$\left(x + \frac{1}{2}\right)^2 - \frac{1}{4}$$



30. a. Less than 4 indicates the parabola faces down

$$\begin{aligned} y &= -(x+1)(x-3) = -(x^2 - 2x - 3) = -(x^2 - 2x - \quad) + 3 \\ &= -(x^2 - 2x + 1) + 3 + 1 \\ &= -(x^2 - 1)^2 + 4 \end{aligned}$$


30. b.  $y = 2x^2 - 4x + 3$

$$y = 2(x^2 - 2x \quad) + 3$$

$$y = 2(x^2 - 2x + 1) + 3 - 2$$

$$y = 2(x-1)^2 + 1$$

vertex (1,1) goes up 2x faster than  $x^2$



31. a.  $x = t + 1$      $t = x - 1$      $y = (x-1)^2 - (x-1)$

$$y = x^2 - 2x + 1 - x + 1$$

$$y = x^2 - 3x + 2$$


31. b.  $x = \sqrt[3]{t} - 1$      $\sqrt[3]{t} = x + 1$      $t = (x+1)^3$      $y = (x+1)^3 [(x+1)^3 - 1]$  Acceptable Answer

$$y = (x+1)^3 [x^3 + 3x^2 + 3x + 1 - 1]$$

$$y = (x+1)^3 [x^3 + 3x^2 + 3x]$$

$$y = x(x+1)^3 [x^2 + 3x + 3]$$

31. c.  $x = \sin t$      $t = \sin^{-1} x$      $y = \cos(\sin^{-1} x)$   
 $y = \sqrt{1 - x^2}$   
 $y^2 = 1 - x^2$   
 $x^2 + y^2 = 1$




32. a.  $x = 2y + 3$   
 $2y = x - 3$   
 $y = \frac{x-3}{2}$


32. b.  $x = \frac{y+2}{5y-1}$   
 $(5y-1)x = y+2$   
 $5yx - x = y+2$   
 $5yx - y = x+2$   
 $y(5x-1) = x+2$   
 $y = \frac{x+2}{5x-1}$

32. c.  $x = y^2 + 2y - 1$   
 $0 = y^2 + 2y - 1 - x$   
 $0 = y^2 + 2y - (1+x)$      $a=1, b=2, c=-(1+x)$   
 $y = \frac{-2 \pm \sqrt{4 - 4(-(1+x))}}{2} = \frac{-2 \pm \sqrt{4+4+4x}}{2} = \frac{-2 \pm \sqrt{8+4x}}{2} = \frac{-2 \pm \sqrt{4(2+x)}}{2} = -1 \pm \sqrt{2+x}$   
 When  $x=0$  in the original equation  $y=-1$ , so  $x$  is limited to  $x > -1$  for the inverse.

33. reflects across  $y = x$




34. a.



Similar Triangles  $\frac{x+t}{t} = \frac{r}{h}$   
 $x+t = \frac{rt}{h}$   
 $x = \frac{rt}{h} - t$   
 $x = \frac{rt}{h} - \frac{th}{h}$   
 $x = \frac{t(r-h)}{h}$

b.




Similar Triangles  $\frac{x}{r} = \frac{t}{\sqrt{r^2 - h^2}}$   
 $x = \frac{rt}{\sqrt{r^2 - h^2}}$


35. a. Area of the square  $= (2r)^2 = 4r^2$  Area of the square  $= \pi r^2$

Area inside the square but outside the circle  $= 4r^2 - \pi r^2$  ratio  $= \frac{4r^2 - \pi r^2}{4r^2} = \frac{4 - \pi}{4} = 1 - \frac{\pi}{4}$


35. b. perimeter  $= r + 2r + r + \frac{2\pi r}{2} = 4r + \pi r$



35. c. Area  $= \pi (1.5)^2 = 2.25\pi$



35. d. Distance  $= \sqrt{100^2 + 200^2} = \sqrt{100^2 + 2^2(100^2)} + \sqrt{(1+4)(100^2)} = 100\sqrt{5}$  km



35. e.  $\sin \theta = \frac{100}{200} \Rightarrow \theta = \sin^{-1} \left( \frac{100}{200} \right) = \frac{\pi}{6}$  or  $30^\circ$